



Contents lists available at ScienceDirect

Planetary and Space Science

journal homepage: www.elsevier.com/locate/pss

Ceres Survey Atlas derived from Dawn Framing Camera images

Th. Roatsch^{a,*}, E. Kersten^a, K.-D. Matz^a, F. Preusker^a, F. Scholten^a, R. Jaumann^a, C.A. Raymond^b, C.T. Russell^c^a Institute of Planetary Research, German Aerospace Center (DLR), Berlin, Germany^b Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA^c Institute of Geophysics, UCLA, Los Angeles, CA, USA

ARTICLE INFO

Article history:

Received 10 November 2015

Received in revised form

26 November 2015

Accepted 16 December 2015

Keywords:

Dawn

Ceres

Dwarf planets

Planetary mapping

ABSTRACT

The Dawn Framing Camera (FC) acquired almost 900 clear filter images of Ceres with a resolution of about 400 m/pixels during the seven cycles in the Survey orbit in June 2015. We ortho-rectified 42 images from the third cycle and produced a global, high-resolution, controlled mosaic of Ceres. This global mosaic is the basis for a high-resolution Ceres atlas that consists of 3 tiles mapped at a scale of 1:2,000,000. The nomenclature used in this atlas was proposed by the Dawn team and was approved by the International Astronomical Union (IAU). The whole atlas is available to the public through the Dawn GIS web page [http://dawn_gis.dlr.de/atlas].

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

The Dawn mission has mapped Ceres during the Survey orbit phase from an altitude of 4424 km (Russell and Raymond, 2011) in seven different cycles. The Dawn spacecraft is equipped with a framing camera [FC (Sierks et al., 2011)] which has one clear filter and seven band pass filters. About 900 clear filter images were taken during Survey, which resulted in a multiple global coverage. We selected 42 images from the third cycle with similar illumination conditions to produce a global mosaic of the illuminated part of Ceres with a resolution of 400 m/pixel, along with a 3-tile atlas at a scale of 1:2,000,000. We describe the different processing steps towards the Ceres Survey atlas, i.e. the image processing steps (Section 2), the Ceres longitude system (Section 3), and finally the cartographic processing aspects (Section 4). We also show the map tiles of the atlas.

2. Data processing

The image data returned from the spacecraft are distributed inside the Dawn team in Planetary Data System (PDS) format [<http://pds.jpl.nasa.gov>]. The first step of the image processing pipeline is the conversion to Video Image Communication and Retrieval (VICAR) format [<http://rushmore.jpl.nasa.gov/vicar.html>]

followed by the radiometric calibration of the images (Schröder et al., 2013,2014).

The next step towards the cartographic products is to ortho-rectify the images to the proper scale and map projection type. This process requires detailed information of the Dawn orbit and attitude data and of the topography of the targets. Both improved orientation and a high-resolution shape model are provided by stereo processing (bundle block adjustment) of the Survey stereo image dataset (Preusker et al., 2015), Ceres's shape model is used for the calculation of the ray intersection points, while the map projection itself is placed onto a reference sphere of Ceres with a mean radius of 470 km. The final step is the mosaicking of all images to a global mosaic of Ceres, the so-called basemap.

The Dawn team proposed 61 names for geological features. By international agreement, craters must be named after gods and goddesses of agriculture and vegetation from world mythology, whereas other geological features must be named after agricultural festivals of the world. The nomenclature proposed by the Dawn team was approved by the IAU [<http://planetarynames.wr.usgs.gov/>] and is shown in Fig. 1.

3. Longitude system

The Ceres longitude system was defined by Feature #1 from the Hubble Space Telescope (HST) mosaic in Li et al. (2006). This feature defined the zero-longitude meridian (Archinal et al., 2011). We could not clearly define this feature in our data but found good agreement for other features, e.g., Feature #2 is close to crater Dantu within the errors

* Corresponding author.

E-mail address: Thomas.Roatsch@dlr.de (Th. Roatsch).




Fig. 1. Global Survey mosaic with nomenclature in Mollweide projection.




Fig. 2. Crater Kait in Survey image FC21A0038023. The upper left box is overlaid by a grid with one degree intervals in latitude and longitude direction.

of the HST images. Therefore, we proposed to the IAU Working Group on Cartographic Coordinates and Rotational Elements (WGCCRE) to select a tiny crater which is close to the former zero degree longitude as a new reference crater. This proposal was accepted, and we selected crater Kait (Fig. 2) as the new reference crater. This crater is now at zero degrees longitude and will be kept at this longitude also in higher resolution mosaics.

4. Ceres map tiles

The Ceres atlas consists of 3 map tiles in the subdivision of the synoptic format. The tiling scheme conforms to those proposed by

Greeley and Batson (1990) and is used e.g., for mapping the Saturnian moons Iapetus and Mimas on a scale of 1:3,000,000 and 1:1,000,000, respectively (Roatsch et al., 2009, 2013). A map scale of about 1:2,000,000 guarantees a mapping at the highest available Dawn FC Survey resolution and results in an acceptable printing scale for the hardcopy map of 5 pixel/mm. The individual map tiles were extracted from the global mosaic and reprojected, coordinate grids were superposed as graphic vectors, and the resulting composites were converted to the common PDF-format. The equatorial map tiles (from 57°S to 57°N latitude) are in Mercator projection with true scale at 0° latitude, whereas the poles are in stereographic projection with true scale at $\pm 90^\circ$ latitude (from 90°S to 55°S latitude and from 55°N to 90°N latitude) (Snyder, 1987). Individual tiles have no overlap in the




Fig. 3. Survey atlas schema.




Fig. 4. Survey atlas quadrangle 1/4 (Asari-Zadeni).




Fig. 5. Survey atlas quadrangle 2 (Kerwan).




Fig. 6. Survey atlas quadrangle 3 (Occator).

east–west direction (see Fig. 3). The longitudinal extension of the map tiles is 180° for those in the Mercator projection and 360° for the pole caps. Using this tiling scheme for Ceres with a high-resolution mosaic, the printed map sheets of the Mercator projections have a size of 110 cm in width by 93 cm in height, and the map sheet of the poles has a size of 110 cm in width by 91 cm in height. Contour lines were derived from a digital terrain model (DTM) of Ceres (Preusker et al., 2015) featuring a lateral resolution of 410 m/pixel with a 3D control point accuracy of ± 60 m. The depicted heights are geometric heights and refer to an oblate ellipsoid with a semi-major axis of 482 km and a semi-minor axis of 446 km. Topography maps showing the terrain with color-coded heights and image index maps were added to every individual tile. The map tiles of the atlas are shown in Figs. 4–6.

The entire Ceres Survey atlas is available to the public through the Dawn GIS web page [http://dawn_gis.dlr.de/atlas].

5. Outlook

The Dawn spacecraft has been in the High Altitude Mapping Orbit (HAMO) since August 18, 2015, and acquires images with a resolution of about 140 m/pixel. Global coverage will give us the

opportunity to calculate a higher resolution atlas of Ceres, similar to the HAMO atlas of Vesta (Roatsch et al., 2012).

Acknowledgments

The authors gratefully acknowledge helpful discussions with B. Archinal (USGS) about the definition of the longitude system and with R. Hayward (USGS) about the proposed nomenclature for Ceres features.

References

- Archinal, B.A., A'Hearn, M.F., Bowell, E., Conrad, A., Consolmagna, G.J., Courtin, R., Fukushima, T., Hestroffer, D., Hilton, J.L., Krasimsky, G.A., Neumann, G., Oberst, J., Seidelmann, P.K., Stooke, P., Tholen, D.J., Thomas, P.C., Williams, L.P., 2011. Report of the IAU Working Group on Cartographic Coordinates and Rotational Elements: 2009. *Celest. Mech. Dyn. Astron.* 109, 101–135.
- Greeley, R., Batson, G., 1990. Planetary Mapping. Cambridge University Press.
- Li, J.-Y., McFadden, L.A., Parker, J. Wm., Young, E.E., Stern, S.A., Thomas, P.C., Russell, C.T., Sykes, M.V., 2006. Photometric analysis of 1 Ceres and surface mapping from HST observations. *Icarus* 182, 143–160.
- Preusker, F., Scholten, F., Matz, K.-D., Roatsch, T., Elgner, S., Jaumann, R., Joy, S.P., Polansky, C.A., Raymond, M.D., Raymond, C.A., Russell, C.T., 2015. Shape model

- and rotational state of dwarf planet Ceres from Dawn FC stereo images. EPSC Meeting. Abstract 186.
- Roatsch, Th., Wählisch, M., Hoffmeister, A., Kersten, E., Matz, K.-D., Scholten, F., Wagner, R., Denk, T., Neukum, G., Porco, C.C., 2009. High-resolution atlases of Mimas, Tethys, and Iapetus derived from Cassini-ISS images. *Planet. Space Sci.* 57, 83–92.
- Roatsch, Th., Kersten, E., Matz, K.-D., Preusker, F., Scholten, F., Jaumann, R., Raymond, C.A., Russell, C.T., 2012. High resolution Vesta High Altitude Mapping Orbit Atlas derived from Dawn Framing Camera images. *Planet. Space Sci.* 73, 283–286.
- Roatsch, Th., Kersten, E., Hoffmeister, A., Wählisch, M., Matz, K.-D., Porco, C.C., 2013. Recent improvements of the Saturnian satellite atlases: Mimas, Enceladus, and Dione. *Planet. Space Sci.* 77, 118–125.
- Russell, C.T., Raymond, C.A., 2011. The Dawn Mission to Vesta and Ceres. *Space Sci. Rev.* 163, 3–23.
- Schröder, S.E., Maue, T., GutiérrezMarqués, P., Mottola, S., Aye, K.M., Sierks, H., Keller, H.U., Nathues, A., 2013. In-flight calibration of the Dawn Framing Camera. *Icarus* 226, 1304–1317.
- Schröder, S.E., Mottola, S., Matz, K.-D., Roatsch, Th., 2014. In-flight calibration of the Dawn Framing Camera II: flat fields and stray light correction. *Icarus* 234, 99–108.
- Sierks, H., Keller, H.-U., Jaumann, R., Michalik, H., Behnke, T., Bubenhausen, F., Büttner, I., Carsenty, U., Christensen, U., Enge, R., Fiethe, B., Gutierrez Marques, P., Hartwig, H., Krüger, H., Kühne, W., Maue, T., Mottola, S., Nathues, A., Reiche, K.-U., Richards, M.L., Roatsch, T., Schröder, S.E., Szemerey, I., Tschentscher, M., 2011. The Dawn Framing Camera. *Space Sci. Rev.* 163, 263–327.
- Snyder, J.P., 1987. *Map Projections – A Working Manual*. US Government Printing Office, Washington, p. 42.