

Hawaii Kuiper Belt Variability Project: An Update

Scott S. Sheppard and David C. Jewitt

Institute for Astronomy, University of Hawaii, 2680 Woodlawn Dr., Honolulu, HI 96822

Abstract.

We have been systematically monitoring a large sample of bright Kuiper Belt objects for possible light variations due to rotational and phase angle effects. Here we report on three objects, 2003 AZ₈₄, (24835) 1995 SM₅₅ and (55636) 2002 TX₃₀₀ observed to have measurable rotational lightcurves with peak-to-peak amplitudes of 0.14 ± 0.03 , 0.19 ± 0.05 and 0.08 ± 0.02 magnitudes and single-peaked periods of 6.71 ± 0.05 , 4.04 ± 0.03 and 8.12 ± 0.08 hours respectively. We observed a further ten objects which showed no rotational photometric variation within measurement uncertainties. In addition, we find that the lightcurve of 1995 SM₅₅ may have a variable amplitude. We discuss this peculiar object as well as our observations of the reportedly variable Kuiper Belt object (19308) 1996 TO₆₆. Finally, we continue to find the phase functions of the Kuiper Belt objects to be very steep and linear, to first order, with a median slope of 0.16 ± 0.01 magnitudes per degree in the phase angle range 0 to 2 degrees.

1. Introduction

The rotations and shapes of the KBOs are probably a function of their size. KBOs may be structurally weak bodies held together by gravity in a rubble pile type structure (Jewitt and Sheppard 2002). The spins of the larger objects are probably primordial with little modification by post-formation impacts. The smaller objects are probably collisional fragments with sizes, shapes and spins determined at the moment of catastrophic break-up (Farinella & Davis 1996). The vast majority of Kuiper Belt Objects (KBOs) currently can not be resolved at their large heliocentric distances ($\gtrsim 30$ AU). Presently, the only feasible way to determine KBO shapes and surface features is by observing their light variations.

We are obtaining voluminous time resolved optical photometric observations to determine the rotational lightcurves and phase functions of KBOs. The University of Hawaii 2.2 meter telescope with its Tektronix 2048×2048 CCD was used for the observations (see section 2 of Sheppard and Jewitt (2002) for further observational and data reduction details). As our sample, we select the intrinsically brightest (presumably largest) KBOs. The 33 large KBOs (≥ 200 km in diameter, assuming a low albedo) we have observed already exceed the number of asteroids of this size.



© 2003 Kluwer Academic Publishers. Printed in the Netherlands.

This short note is a continuation of Jewitt and Sheppard (2002) and Sheppard and Jewitt (2002) and a more detailed write-up will soon follow. The present work is intended as a timely pointer to a manuscript that is in preparation and our already published articles and is not a substitute for them.

2. Rotation Results

Rotational lightcurve analyses for the newly observed KBOs are shown in Table 1 and discussed below. We employed the phase dispersion minimization (PDM) method (Stellingwerf 1978) to search for periodicity in the data. The best-fit period should have a very small normalized dispersion, Θ , compared with the unphased data, and thus $\Theta \ll 1$ indicates that a good fit has been found.

When combining the newly observed objects with our previous data (Sheppard and Jewitt 2002) we find that 9 of 33 (27%) objects in our sample show peak-to-peak amplitudes ≥ 0.15 magnitudes while 15% have amplitudes ≥ 0.4 magnitudes and 9% have amplitudes ≥ 0.6 magnitudes. The large main-belt asteroids have a larger fraction of objects with amplitudes ≥ 0.15 magnitudes, a comparable fraction with amplitudes ≥ 0.4 magnitudes but a smaller fraction with amplitudes ≥ 0.6 magnitudes.

Please see Jewitt and Sheppard (2002) as well as Sheppard and Jewitt (2002) for further discussion on the interpretation of KBO lightcurves. The complete data from these new results will soon be published along with further observations in Sheppard and Jewitt (2003).

2.1. 2003 AZ₈₄

PDM analysis shows that 2003 AZ₈₄ has a single-peaked lightcurve period of $P = 6.72 \pm 0.05$ hours (Figure 1). One minimum and one maximum in brightness within a single night were observed and put the full single-peaked lightcurve just over 6 hours, in agreement with the PDM analysis. We phased the data and found the 6.72 hour period to be very good (Figure 2). The peak-to-peak variation is $\Delta m = 0.14 \pm 0.03$ magnitudes. We possess no evidence to show whether the lightcurve is singly periodic (as expected from surface albedo variations) or doubly periodic (consistent with aspherical shape).

2.2. (24835) 1995 SM₅₅

We observed (24835) 1995 SM₅₅ for five nights in October 2001. The KBO was found to have very scattered photometry for its brightness (error bars on the photometry are only 0.03 magnitudes), and a good periodic lightcurve could not be identified. We thus reported this object as having a flat lightcurve in our secondary sample in Sheppard and Jewitt (2002). Further observations were obtained in November 2001 to determine if the scattered photometry was caused by noise. With the additional photometry we were able to find a rotational lightcurve with single-peaked period near $P = 4$ hours (Figure 3). Phasing the data together again showed that the single-peaked lightcurve was extremely noisy given our uncertainties (Figure 4). Phasing the data to a possible double-peaked lightcurve of 8 hours is also very noisy (Figure 5). An average peak-to-peak lightcurve amplitude is 0.19 ± 0.05 . It appears that the amplitude of the rotational lightcurve may be variable from night to night.

Trujillo and Brown (2003) observed 1995 SM₅₅ with the Hubble Space Telescope and found no evidence that it is a binary object. However, their constraint applies only to satellites with separation ≥ 0.1 arcseconds and having a magnitude difference ≤ 2.5 . Therefore it remains possible that the “noisy” lightcurve of 1995 SM₅₅ is due to the presence of multiple periods in the photometric data. Only protracted, highly accurate photometric series can show whether or not this is the case. It is interesting that 1995 SM₅₅ is one of the bluest KBOs known ($V-R = 0.38$; see Hainaut & Delsanti 2002). It may be that blue objects have recently had their volatile rich insides exposed, possibly by a recent collision. Thus the amplitude variation seen for 1995 SM₅₅ may be affected by cometary activity from this freshly exposed material. The large scatter in photometry could also be due to the object being in a complex rotational state, although it is difficult to see how such a state could be maintained other than by forced precession due to a satellite or a very unlikely recent collision since the damping time for wobbles is short (Burns & Safronov 1973). Further observations of 1995 SM₅₅ are needed to understand its peculiar lightcurve nature.

2.3. (19308) 1996 TO₆₆

A similar situation has been invoked for the very blue object (19308) 1996 TO₆₆ in which the lightcurve was reported to show signs of variability (Hainaut et al. 2000; Sekiguchi et al. 2002). We observed 1996 TO₆₆ in November 2001 and PDM analysis of our data shows there are possible rotational single-peaked lightcurves at about 3.96 ± 0.04 and 4.80 ± 0.05 hours and double-peaked periods around 5.90 ± 0.05 ,

7.92 ± 0.04 and 9.6 ± 0.1 hours (Figure 6). When phasing the data to these values, all seem plausible, but the single-peaked period near 3.96 hours and the corresponding double-peaked period near 7.92 hours are significantly better (Figures 7 and 8). We find the peak-to-peak amplitude of 1996 TO₆₆ to be 0.26 ± 0.03 magnitudes in our 2001 observations.

Hainaut et al. (2000) reported a 6.25 hour period for 1996 TO₆₆ in data taken in 1997 (0.12 magnitudes in amplitude) and 1998 (0.33 magnitudes) and later affirmed this period in photometry from 1999 (0.21 magnitudes; Sekiguchi et al. 2002). They suggest that changes in the lightcurve period from double-peaked to single-peaked as well as in the amplitude between 1997 and 1998 can be attributed to activity, possibly outgassing, in 1996 TO₆₆. Phasing our 2001 data to 6.25 hours gives an implausible lightcurve. We found that the very sparsely sampled Sekiguchi et al. (2002) data from 1999 observations of 1996 TO₆₆ are consistent with many periods, including the periods found in our 2001 data. O. Hainaut kindly provided us with the photometric measurements for the 1997 and 1998 observations of 1996 TO₆₆ described in Hainaut et al. (2000); the data currently can also be found on the Small Bodies Node of the Planetary Data System at <http://www.psi.edu/pds/tnolc.html>. In examining this data we could not find a significant lightcurve in the 1997 data (≤ 0.1 magnitudes in amplitude). In our PDM analysis of the Hainaut et al. data from 1998 we found similar periods for 1996 TO₆₆ as our 2001 observations (Figure 9). We do not find strong evidence for the 6.25 hour period in the Hainaut et al. data from 1998.

We do not see any evidence that the period of 1996 TO₆₆ has changed between the 1997, 1998, 1999 and 2001 observations. Romanishin and Tegler (1999) found the lightcurve for 1996 TO₆₆ in 1997 to be ≤ 0.1 magnitudes, seemingly corroborating the small amplitude reported by Hainaut et al. Thus, while there is no evidence for a change in rotation period, the lightcurve amplitude of 1996 TO₆₆ may have changed since 1997.

2.4. (55636) 2002 TX₃₀₀

PDM analysis shows that (55636) 2002 TX₃₀₀ has a single-peaked lightcurve period of either $P = 8.12 \pm 0.08$ or $P = 12.10 \pm 0.08$ hours (Figure 10). Both single-peaked periods appear acceptable in the phased data (Figure 11) with a peak-to-peak variation of $\Delta m = 0.08 \pm 0.02$ magnitudes.

2.5. FLAT ROTATIONAL LIGHTCURVE OBJECTS

Ten KBOs ((55637) 2002 UX₂₅, (55638) 2002 VE₉₅, (47171) 1999 TC₃₆, (42355) 2002 CR₄₆, (28978) Ixion 2001 KX₇₆, 2000 YW₁₃₄, (42301) 2001 UR₁₆₃, 2001 QF₂₉₈, 2001 FP₁₈₅, and 2001 KD₇₇) showed no measurable photometric variations (Table 1), by which we mean that their lightcurves have range ≤ 0.1 magnitudes and/or period > 24 hours. A few objects show hints of variability that might, with more data, emerge as rotationally modulated lightcurves. The KBO 2001 YW₁₃₄ has a variation of about 0.1 magnitudes near a 5 hour single-peaked period on one night, but the object appears mostly flat on the second night over 5 hours. Finally, the faint KBO 2001 QF₂₉₈ appears to have variations of about 0.1 magnitudes. Confirmation of these subtle lightcurves will require more data, with a larger telescope most likely required.

3. Phase Angle Results

We add additional measurements for four KBO phase functions reported in Sheppard and Jewitt (2002) as well as six new KBOs (Table 2). We continue to find that the slopes are very steep and the additional points show that the phase functions are linear to first order between phase angles of 0 and 2 degrees (Figure 12). For the phase functions we use the notation $\phi(\alpha) = 10^{-0.4\beta\alpha}$, where α is the phase angle in degrees and β is the “linear” phase coefficient. The median phase coefficient using all 13 KBOs observed at significantly different phase angles from Sheppard and Jewitt (2002) and this work is $\beta = 0.16 \pm 0.01$ magnitudes per degree. Though not necessarily useful at the low phase angles for which KBOs can be observed, we also include the H and G formalism as described in Bowell et al. (1989) in Table 2 in order to fully compare our results with other works. As previously, we note that the large, high albedo object Pluto has a much lower phase function than the smaller KBOs (0.0294 ± 0.0011 mag/deg; Buie, Tholen and Wasserman 1997). The intermediate albedo/sized Charon has an intermediate phase function (0.0866 ± 0.0078 mag/deg; Buie, Tholen and Wasserman 1997). Although there exists no unique correlation between albedo and phase function at low phase angles, our data are consistent with low albedos for the 100-1000 km scale KBOs.

Acknowledgements

This work was supported by a grant to D.J. from the NASA Origins Program.

References

- Burns, J. and Safronov, V. Asteroid Nutation Angles. *MNRAS*, 165:403–411, 1973.
- Bowell, E. et al. Application of photometric models to asteroids. In R. Binzel, T. Gehrels and M. Matthews, editors, *Asteroids II*, Tucson, Arizona, 1989
- Buie, M., Tholen, D. and Wasserman, L. Separate Lightcurves of Pluto and Charon. *Icarus*, 125:233–244, 1997.
- Farinella, P. and Davis, D. Short-period comets: Primordial bodies or collisional fragments? *Science*, 273:938–941, 1996.
- Hainaut, O. et al. Physical Properties of TNO 1996 TO66. Lightcurves and possible cometary activity. *A&A*, 356:1076–1088, 2000.
- Hainaut, O. and Delsanti, A. Colors of Minor Bodies in the Outer Solar System: A statistical Analysis. *A&A*, 389:641–664, 2002.
- Jewitt, D. and Sheppard, S. Physical Properties of Trans-Neptunian Object (20000) Varuna. *Astron. J.*, 123:2110–2120, 2002.
- Romanishin, W. and Tegler, S. Rotation rates of Kuiper-belt objects from their light curves. *Nature*, 398:129–132, 1999.
- Sekiguchi, T., Boehnhardt, H., Hainaut, O. and Delahodde, C. Bicolour Lightcurve of TNO 1996 TO66 with the ESO-VLT. *A&A*, 385:281–288, 2002.
- Sheppard, S. and Jewitt, D. Time-Resolved Photometry of Kuiper Belt Objects: Rotations, Shapes, and Phase Functions. *Astron. J.*, 124:1757–1775, 2002.
- Sheppard, S. and Jewitt, D. in preparation 2003.
- Stellingwerf, R. Period Determination Using Phase Dispersion Minimization. *Astron. J.*, 224:953–960, 1978.
- Trujillo, C. and M. Brown A Search for Bright Kuiper Belt Binaries with the Hubble Space Telescope. *Astron. J.* submitted, 2003

Figure 1. The phase dispersion minimization (PDM) plot for 2003 AZ₈₄. Best fits from this plot are the 6.72 hour single-peaked fit and the 13.44 hour double-peaked fit.

Figure 2. Phased R-band data from the UT February 2003 observations for 2003 AZ₈₄. The period has been phased to the single-peaked period of 6.72 hours. Uncertainties for individual points are ± 0.03 .

Figure 3. The PDM plot for 1995 SM₅₅. Best fits from this plot are the 4.04 hour single-peaked fit and the 8.08 hour double-peaked fit. Both are flanked by aliases.

Figure 4. Single-peaked 4.04 hour phased R-band data from the UT October and November 2001 observations for 1995 SM₅₅. Uncertainties for individual points are ± 0.03 .

Figure 5. Double-peaked 8.08 hour phased R-band data from the UT October and November 2001 observations for 1995 SM₅₅. Uncertainties for individual points are ± 0.03 .

Figure 6. The PDM plot for 1996 TO₆₆. Best fits from this plot are the 3.96 hour single-peaked fit and the 7.92 hour double-peaked fit.

Figure 7. Single-peaked 3.96 hour phased R-band data from the UT October 2001 observations for 1996 TO₆₆. Uncertainties for individual points are ± 0.03 .

Figure 8. Double-peaked 7.92 hour phased R-band data from the UT October 2001 observations for 1996 TO₆₆. Uncertainties for individual points are ± 0.03 .

Figure 9. The PDM plot for 1996 TO₆₆ using the observations in 1998 taken by Hainaut et al. (2000). This PDM plot looks very similar to our 2000 observations PDM plot with good 3.96 hour single-peaked and 7.92 hour double-peaked fits.

Figure 10. The phase dispersion minimization (PDM) plot for 2002 TX₃₀₀. Best fits from this plot are the 8.12 and 12.10 hour single-peaked fits.

Figure 11. Phased R-band data from the UT November 2002 observations for 2002 TX₃₀₀. The period has been phased to the single-peaked period of 8.12 hours. Uncertainties for individual points are ± 0.02 .

Figure 12. Phase functions for the new observations of KBOs observed at several phase angles. The KBO phase functions appear linear to first order. Reduced magnitudes have been normalized and offset in order to display the phase functions efficiently.

TABLE 1. Properties of Newly Observed KBOs

Name	m_R^a (mag)	Nights ^b (#)	Δm_R^c (mag)	Single ^d (hrs)	Double ^e (hrs)	
(55636)	2002 TX ₃₀₀	19.29 ± 0.04	3	0.08 ± 0.02	8.12 ± 0.08 12.10 ± 0.08	16.24 ± 0.08 24.20 ± 0.08
(55637)	2002 UX ₂₅	19.65 ± 0.02	2	< 0.06	-	-
(55638)	2002 VE ₉₅	19.68 ± 0.02	1	< 0.06	-	-
(47171)	1999 TC ₃₆	19.80 ± 0.02	2	< 0.05	-	-
(42355)	2002 CR ₄₆	19.82 ± 0.02	4	< 0.05	-	-
(28978) Ixion	2001 KX ₇₆ ^f	19.84 ± 0.02	3	< 0.05	-	-
	2003 AZ ₈₄	20.14 ± 0.07	3	0.14 ± 0.03	6.72 ± 0.05	13.44 ± 0.05
(24835)	1995 SM ₅₅ ^f	20.20 ± 0.10	9	0.19 ± 0.05	4.04 ± 0.03	8.08 ± 0.03
	2000 YW ₁₃₄	20.67 ± 0.05	2	< 0.1	?	?
(42301)	2001 UR ₁₆₃	20.86 ± 0.03	3	< 0.08	-	-
(19308)	1996 TO ₆₆ ^f	21.12 ± 0.13	5	0.26 ± 0.03	3.96 ± 0.04	7.92 ± 0.04
	2001 FP ₁₈₅	21.15 ± 0.03	4	< 0.06	-	-
	2001 QF ₂₉₈	21.41 ± 0.06	4	< 0.12	?	?
	2001 KD ₇₇	21.48 ± 0.03	3	< 0.07	-	-

^aMean R-band magnitude on the date having the majority of observations.

^bNumber of nights used to determine the lightcurve.

^cThe peak to peak range of the lightcurve.

^dThe lightcurve period if there is one maximum per period.

^eThe lightcurve period if there is two maxima per period.


^fInitial details were given in our secondary sample in Sheppard and Jewitt (2002). Additional observations have been obtained and thus we give further details on the objects here.


TABLE 2. New Phase Function Data for KBOs


Name	H	G	$\beta(\alpha < 2^\circ)^a$	N^b
(42301) 2001 UR163	3.75 ± 0.05	-1.00 ± 0.30	0.25 ± 0.05	2
2000 YW134	4.29 ± 0.20	$+0.57 \pm 0.50$	0.07 ± 0.12	2
(38628) Huya 2000 EB173	4.45 ± 0.01	$+0.10 \pm 0.03$	0.11 ± 0.01	4
(26181) 1996 GQ21	4.47 ± 0.01	$+0.12 \pm 0.03$	0.12 ± 0.02	4
(26375) 1999 DE9	4.55 ± 0.02	-0.34 ± 0.05	0.17 ± 0.05	4
(26375) 1997 CS29	4.78 ± 0.10	-1.42 ± 0.30	0.31 ± 0.10	3
2001 QF298	4.95 ± 0.10	-0.19 ± 0.10	0.16 ± 0.03	2
2001 CZ31	5.54 ± 0.02	-0.03 ± 0.05	0.13 ± 0.03	3
2001 FZ173	5.63 ± 0.03	-0.77 ± 0.08	0.23 ± 0.03	4
2001 FP185	5.79 ± 0.05	-1.13 ± 0.20	0.28 ± 0.05	2


^a $\beta(\alpha < 2^\circ)$ is the phase coefficient at phase angles $< 2^\circ$.


^b N is the number of different observing runs used and thus the number of points plotted. Most objects were observed on many consecutive nights during each of several observing runs. These data were all averaged together to obtain points with very small uncertainties.





























Normalized Reduced R-Magnitude

